



ACETAMINOPHEN DOSING: Doses to be given a minimum of 4 hours apart

WEIGHT	INFANT DROPS * 80 MG/0.8 ML	SUSPENSION/LIQUID * 160 MG/5 ML	CHEWABLE TABLET 80 MG	CHEWABLE TABLET 160 MG	ADULT TABLET 325 MG
6-11 lb	½ dropper = 0.4 ml	¼ teaspoon = 1.25 ml			
12-17 lb	1 dropper = 0.8 ml	½ teaspoon = 2.5 ml			
18-23 lb	1 ½ dropper = 1.2 ml	¾ teaspoon = 3.75 ml			
24-35 lb	2 dropper = 1.6 ml	1 teaspoon = 5 ml	2 tablets	1 tablet	
36-47 lb		1 ½ teaspoons = 7.5 ml	3 tablets	1 ½ tablets	
48-59 lb		2 teaspoons = 10 ml	4 tablets	2 tablets	1 tablet
60-71 lb		2 ½ teaspoons = 12.5 ml	5 tablets	2 ½ tablets	1 tablet
72-95 lb		3 teaspoons = 15 ml	6 tablets	3 tablets	1 ½ tablets
> 95 lb		4 teaspoons = 20 ml	8 tablets	4 tablets	2 tablets

*Please note that there are both infant drops (80 mg/0.8 ml) and infant suspension (160 mg/5 ml) and dose accordingly

IBUPROFEN DOSING:

- Doses to be given a minimum of 6 hours apart; not recommended for children under 6 months old

WEIGHT	INFANT DROPS 50 MG/ 1.25ML	LIQUID 100 MG/ 5 ML	CHEWABLE TABLET 50 MG	CHEWABLE TABLET 100 MG	ADULT TABLET 200 MG
12-17 lb	1 dropper = 1.25 ml	½ teaspoon = 2.5 ml			
18-23 lb	1½ dropper = 1.875 ml	¾ teaspoon = 3.75 ml			
24-35 lb	2 dropper = 2.5 ml	1 teaspoon = 5 ml	2 tablets	1 tablet	
36-47 lb		1 ½ teaspoons = 7.5 ml	3 tablets	1 ½ tablets	
48-59 lb		2 teaspoons = 10 ml	4 tablets	2 tablets	1 tablet
60-71 lb		2 ½ teaspoons = 12.5 ml	5 tablets	2 ½ tablets	1 tablet
72-92 lb		3 teaspoons = 15 ml	6 tablets	3 tablets	1 ½ tablets
> 92 lb		4 teaspoons = 20 ml	8 tablets	4 tablets	2 tablets

BENADRYL (Diphenhydramine): Doses to be given every 4-6 hours as needed

Weight	Elixir 12.5 mg/ 5ml	Chewable Tablets 12.5 mg	Tablets 25 mg
12-17 lb	½ teaspoon = 2.5 ml		
18-23 lb	¾ teaspoon = 3.75 ml		
24-35 lb	1 teaspoon = 5 ml	1 tablet	
36-47 lb	1 ½ teaspoons = 7.5 ml	1 ½ tablets	
48-59 lb	2 teaspoons = 10 ml	2 tablets	1 tablet
60-71 lb		2 tablets	1 tablet
72-95 lb		2 tablets	1 tablet
> 95 lb		2 tablets	1 tablet